

SN 10/646,378
Docket No. S-94,729
In Response to Office Action dated September 14, 2005

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS:

1. (Currently amended) A functionalized polymer for binding a solute or a suite of solutes dissolved in a solution comprising:

a polymeric backbone; and

one or more functional groups covalently linked to the polymeric backbone, the one or more functional groups selected to bind to the solute or a suite of solutes, the one or more functional groups selected from the group consisting of a monol derivative, a diol derivative, a triol derivative, a tetraol derivative, a glucarone derivative, a thiol derivative, a dithiol derivative, an alpha-hydroxycarboxylic acid derivative, a tartrate derivative, a calixarene derivative, a polypeptide derivative, bisphosphonic acid derivatives, biscarboxylic acid derivatives and bisamide derivatives, bisester derivatives, a monoester derivative, a monoamide derivative, a mixed phosphonic acid/carboxylic acid derivative, an alkylpyridinium derivative, a cyclodextran derivative, an antibody, a Fab fragment of an antibody, a F(ab)₂ of an antibody, an antigen, a cavity of selected size that hosts the solute selectively, a cage-shaped host, guest-host groups, and affinity groups, wherein the functionalized polymer is bindable to a non-metallic solute or a suite of non-metallic solutes.

2. (Original) The functionalized polymer of claim 1, wherein the polymeric backbone is selected from the group consisting of polyvinylamine, polyallylamine, polyacrylamide, polyethylenimine, polyacrylic acid, polymethacrylic acid, polyvinylalcohol, polyvinylacetate, polypyrrol, and hyperbranched polymers.

3. (Original) The functionalized polymer of claim 1, wherein the diol derivative is tartrate.

4. (Original) The functionalized polymer of claim 3, wherein the tartrate is covalently linked to the polymeric backbone to from a cyclic tartrate imide.

SN 10/646,378

Docket No. S-94,729

In Response to Office Action dated September 14, 2005

5. (Original) The functionalized polymer of claim 3, wherein the tartrate is covalently linked to the 5 polymeric backbone to form an open monoester, which can be converted to a carboxylic acid.

6. (Original) The functionalized polymer of claim 3, wherein the tartrate is covalently linked to the polymeric backbone to form a diamide attached to two different nitrogen atoms of the polymeric backbone.

7. (Original) The functionalized polymer of claim 1, wherein the diol derivative is glycidol.

8. (Original) The functionalized polymer of claim 1, wherein the functionalized polymer is crosslinked thereby rendering the functionalized polymer insoluble in water.

9. (Original) The functionalized polymer of claim 1, wherein the functionalized polymer is water-soluble.

10. (Original) The functionalized polymer of claim 9, wherein the functionalized polymer is purified to have polymer molecule sizes capable of being retained by a membrane with a molecular weight cutoff value of a first pre-selected level and essentially free of polymer molecule sizes capable of passing through a membrane with a molecular weight cutoff value of a second pre-selected level the second pre-selected level being smaller than the first pre-selected level.

11. (Currently amended) The functionalized polymer of claim 10, wherein the functionalized polymer has a molecular weight in the range from about 1,000 MW to about 1,000,000 MW.

12. (Currently amended) The functionalized polymer of claim 10, wherein the functionalized polymer has a molecular weight in the range from about 10,000 MW to about 100,000 MW.

13. (Original) The functionalized polymer of claim 1, wherein the functionalized polymer is represented by the formula X-R wherein "X" is a synthetic polymer selected from the group consisting of polyethylenimine, polyvinylamine, polyallylamine, polypropylamine, polyacrylamide, polyethylenimine, polyacrylic acid, polymethacrylic acid, polyvinylalcohol, polyvinylacetate, polypyrrol, or hyperbranched polymers and "R" is a functional group selected from the group consisting of a monool derivative, a diol derivative, a tetraol derivative, a triol derivative, an alphahydroxycarboxylate derivative, a glucarone derivative, and a dithiol derivative.

SN 10/646,378
Docket No. S-94,729
In Response to Office Action dated September 14, 2005

14. (Original) The functionalized polymer of claim 13, wherein the polymer is water soluble.

15. (Original) The functionalized polymer of claim 14, wherein the functionalized polymer is purified to have polymer molecule sizes capable of being retained by a membrane with a molecular weight cutoff value of a first pre-selected level and essentially free of polymer molecule sizes capable of passing through a membrane with a molecular weight cutoff value of a second pre-selected level the second pre-selected level being smaller than the first pre-selected level.

16. (Currently amended) The functionalized polymer of claim 14, wherein the functionalized polymer has a molecular weight in the range from about 1,000 MW to about 1,000,000 MW.

17. (Currently amended) The functionalized polymer of claim 14, wherein the functionalized polymer has a 5 molecular weight in the range from about 10,000 MW to about 100,000 MW.

18. (Currently amended) The functionalized polymer of claim 13, wherein the non-metallic solute or suite of non-metallic solutes is selected from the group consisting of species of arsenic, ~~barium, cadmium, chromium, mercury, lead, silver,~~ selenium, actinides, lanthanides, ~~copper, nickel, zinc, cobalt,~~ boron, silicon, and antimony.

19. (Original) The functionalized polymer of claim 1, wherein the functionalized polymer is represented by the formula X-R, wherein "X" is a synthetic polymer selected from the group consisting of polyethylenimine, polyvinylamine, polyallylamine, polypropylamine, polyacrylamide, polyethylenimine, polyacrylic acid, polymethacrylic acid, polyvinylalcohol, polyvinylacetate, polypyrrol, or hyperbranched polymers and "R" is a functional group selected from the group consisting of a thiol derivative, a tartrate derivative, a calixarene derivative, a polypeptide derivative, bisphosphonic acid derivatives, biscarboxylic acid derivatives and bisamide derivatives, bisester derivatives, a monoester derivative, a monoamide derivative, a mixed phosphonic acid/carboxylic acid derivative, an alkylpyridinium derivative, a cyclodextran derivative, an antibody, a Fab fragment of an antibody, a F(ab)₂ of an antibody, an antigen, a cavity of selected size that hosts the solute selectively, a cage-shaped host, guest-host groups, and affinity groups.

SN 10/646,378
Docket No. S-94,729
In Response to Office Action dated September 14, 2005

20. (Original) The functionalized polymer of claim 19, wherein the polymer is water soluble.

21. (Original) The functionalized polymer of claim 20, wherein the functionalized polymer is purified to have polymer molecule sizes capable of being retained by a membrane with a molecular weight cutoff value of a first pre-selected level and essentially free of polymer molecule sizes capable of passing through a membrane with a molecular weight cutoff value of a second pre-selected level the second pre-selected level being smaller than the first pre-selected level.

22. (Currently amended) The functionalized polymer of claim 20, wherein the functionalized polymer has a 10 molecular weight in the range from about 1,000 MW to about 1,000,000 MW.

23. (Currently amended) The functionalized polymer of claim 20, wherein the functionalized polymer has a molecular weight in the range from about 10,000 MW to about 100,000 MW.

24. (Currently amended) The functionalized polymer of claim 19, wherein the non-metallic solute or suite of non-metallic solutes is selected from the group consisting of species of arsenic, barium, cadmium, chromium, mercury, lead, silver, selenium, actinides, lanthanides, copper, nickel, zinc, cobalt, boron, silicon, and antimony.

25. (Currently amended) The functionalized polymer of claim 1, wherein the non-metallic solute or suite of non-metallic solutes is selected from the group consisting of species of arsenic, barium, cadmium, chromium, mercury, lead, silver, selenium, actinides, lanthanides, copper, nickel, zinc, cobalt, boron, silicon, iodine, and antimony.

26. (Currently amended) A functionalized synthetic polymer for binding a solute or a suite of solutes dissolved in a solution comprising:

a water-soluble backbone polymer selected from the group consisting of polyethylenimine, polyvinylamine, polyallylamine, polypropylamine, polyacrylamide, polyethylenimine, polyacrylic acid, polymethacrylic acid, polyvinylalcohol, polyvinylacetate, polypyrrol, or hyperbranched polymers; and

one or more functional groups covalently linked to the polymeric backbone, the one or more functional groups selected to bind to the solute or a suite of solutes, the one or more functional groups selected from the group consisting of a monol derivative, a diol derivative, a thiol derivative, a tetraol derivative, a glucarone derivative, a thiol

SN 10/646,378
Docket No. S-94,729
In Response to Office Action dated September 14, 2005

derivative, a dithiol derivative, an alpha-hydroxycarboxylic acid derivative, a tartrate derivative, a calixarene derivative, a polypeptide derivative, bisphosphonic acid derivatives, biscarboxylic acid derivatives and bisamide derivatives, bisester derivatives, a monoester derivative, a monoamide derivative, a mixed phosphonic acid/carboxylic acid derivative, an alkylpyridinium derivative, a cyclodextran derivative, an antibody, a Fab fragment of an antibody, a F(ab)₂ of an antibody, an antigen, a cavity of selected size that hosts the solute selectively, a cage-shaped host, guest-host groups, and affinity groups, the functionalized polymer being purified to have polymer molecule sizes capable of being retained by a membrane with a molecular weight cutoff value of a first pre-selected level and essentially free of polymer molecule sizes capable of passing through a membrane with a molecular weight cutoff value of a second pre-selected level the second pre-selected level being smaller than the first pre-selected level, wherein the functionalized polymer is bindable to a non-metallic solute or a suite of non-metallic solutes.

27. (Original) The functionalized polymer of claim 26, wherein the diol derivative is tartrate.

28. (Original) The functionalized polymer of claim 27, wherein the tartrate is covalently linked to the polymeric backbone to form a cyclic tartrate imide.

29. (Original) The functionalized polymer of claim 27, wherein the tartrate is covalently linked to the polymeric backbone to form an open monoester.

30. (Original) The functionalized polymer of claim 27, wherein the tartrate is covalently linked to the polymeric backbone to form a diamide attached to two different nitrogen atoms of the polymeric backbone.

31. (Currently amended) The functionalized polymer of claim 26, wherein the functionalized polymer has a molecular weight in the range from about 1,000 MW to about 1,000,000 MW.

32. (Currently amended) The functionalized polymer of claim 26, wherein the functionalized polymer has a molecular weight in the range from about 10,000 MW to about 100,000 MW.

33. (Currently amended) The functionalized polymer of claim 32, wherein the non-metallic solute or suite of non-metallic solutes is selected from the group consisting of

SN 10/646,378

Docket No. S-94,729


In Response to Office Action dated September 14, 2005

species of arsenic, ~~barium, cadmium, chromium, mercury, lead, silver~~, selenium, actinides, ~~lanthanides, copper, nickel, zinc, cobalt~~, boron, silicon, and antimony.

34. (Currently amended) The functionalized polymer of claim 26, wherein the solute or suite of solutes is selected from the group consisting of arsenic, barium, ~~cadmium, chromium, mercury, lead, silver~~, selenium, actinides, ~~lanthanides, copper, nickel, zinc, cobalt~~, boron, silicon, iodine, and antimony.

35. (Original) The functionalized synthetic polymer of claim 1, wherein the functionalized polymer is represented by the formula X-R wherein "X" is a polymer selected from the group consisting of polyethylenimine, polyvinylamine, polyallylamine, polypropylamine polyacrylamide, polyethylenimine, polyacrylic acid, polymethacrylic acid, polyvinylalcohol, polyvinylacetate, polypyrrol, or hyperbranched polymer and "R" is a functional group selected from the group consisting of a monol derivative, a diol derivative, a triol derivative, a tetraol derivative, a glucarone derivative, a thiol derivative, a dithiol derivative, an alpha-hydroxycarboxylic acid deravitive, a tartrate derivative, a calixarene derivative, a polypeptide derivative, bisphosphonic acid derivatives, biscarboxylic acid derivatives and bisamide derivatives, bisester derivatives, a monoester derivative, a monoamide derivative, a mixed phosphonic acid/carboxylic acid derivative, an alkylpyridinium derivative, a cyclodextran derivative, an antibody, a Fab fragment of an antibody, a F(ab)₂ of an antibody, an antigen, a cavity of selected size that hosts the solute selectively, a cage-shaped host, guest-host groups, and affinity groups.

36. (Currently amended) A functionalized polymer for binding a solute or a suite of solutes dissolved in a solution, the functionalized polymer comprising a molecule of the formula:



SN 10/646,378
Docket No. S-94,729
In Response to Office Action dated September 14, 2005

wherein n is an integer between about 12 and about 12,000 and R is NH₂ except at one or more positions within the polymer where R is a functional group independently selected from the group consisting of a monol derivative, a diol derivative, a triol derivative, a tetraol derivative, a glucarone derivative, a thiol derivative, a dithiol derivative, an alpha-hydroxycarboxylic acid derivative, a tartrate derivative, a calixarene derivative, a polypeptide derivative, bisphosphonic acid derivatives, biscarboxylic acid derivatives and bisamide derivatives, bisester derivatives, a monoester derivative, a monoamide derivative, a mixed phosphonic acid/carboxylic acid derivative, an alkylpyridinium derivative, a cyclodextran derivative, an antibody, a Fab fragment of an antibody, a F(ab)₂ of an antibody, an antigen, a cavity of selected size that hosts the solute selectively, a cage-shaped host, guest-host groups, and affinity groups, wherein the functionalized polymer is bindable to a non-metallic solute or a suite of non-metallic solutes.

37. (Original) The functionalized polymer of claim 36, wherein the functionalized polymer is crosslinked thereby rendering the functionalized polymer insoluble in water.

38. (Original) The functionalized polymer of claim 36, wherein the functionalized polymer is water-soluble.

39. (Original) The functionalized polymer of claim 38, wherein the functionalized polymer is purified to 20 have polymer molecule sizes capable of being retained by a membrane with a molecular weight cutoff value of a first pre-selected level and essentially free of polymer molecule sizes capable of passing through a membrane with a molecular weight cutoff value of a second pre-selected level the second pre-selected level being smaller than the first pre-selected level.

40. (Currently amended) The functionalized polymer of claim 39, wherein the functionalized polymer has a molecular weight in the range from about 1,000 MW to about 1,000,000 MW.

41. (Currently amended) The functionalized polymer of claim 39, wherein the functionalized polymer has a molecular weight in the range from about 10,000 MW to about 100,000 MW.

42. (Currently amended) The functionalized polymer of claim 39, wherein the the non-metallic solute or suite of non-metallic solutes is selected from the group consisting of

SN 10/646,378

Docket No. S-94,729

In Response to Office Action dated September 14, 2005

species of arsenic, ~~barium, cadmium, chromium, mercury, lead, silver,~~ selenium, actinides, ~~lanthanides, copper, nickel, zinc, cobalt,~~ boron, silicon, and antimony.

43. (Currently amended) The functionalized polymer of claim 36, wherein the solute or suite of solutes is selected from the group consisting of arsenic, ~~barium, cadmium, chromium, mercury, lead, silver,~~ selenium, actinides, ~~lanthanides, copper, nickel, zinc, cobalt,~~ boron, silicon, iodine, and antimony.